

AD _____

Award Number: DAMD17-99-1-9091

TITLE: Diagnosing Breast Cancer Using Protease Fingerprint

PRINCIPAL INVESTIGATOR: Emily Chen, B.S.

CONTRACTING ORGANIZATION: The Burnham Institute
La Jolla, California 92037

REPORT DATE: June 2000

TYPE OF REPORT: Annual Summary

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

DTIC QUALITY INSPECTED 4

20001019 099

REPORT DOCUMENTATION PAGEForm Approved
OMB No. 074-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE June 2000	3. REPORT TYPE AND DATES COVERED Annual Summary (1 Jun 99 - 31 May 00)	
4. TITLE AND SUBTITLE Diagnosing Breast Cancer Using Protease Fingerprint			5. FUNDING NUMBERS DAMD17-99-1-9091	
6. AUTHOR(S) Emily Chen, B.S.				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) The Burnham Institute La Jolla, California 92037 E-MAIL: echen@burnham-ist.org			8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012			10. SPONSORING / MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution unlimited				12b. DISTRIBUTION CODE
13. ABSTRACT (Maximum 200 Words) I constructed a substrate phage display library displaying randomized hexamer amino acid sequences to explore the possibility of profiling protease activity in non-metastatic breast tumor and metastatic breast tumor. This technique was first applied to two breast cancer implicated and closely related metalloproteinase, MMP-2 and MMP-9. Novel substrate specificity was found for MMP-2 vs. MMP-9. These novel substrate sequences can serve as a tool for us to investigate the role of MMP-2 and MMP-9 during breast tumor progression. Moreover, these unique substrate sequences can be used as leads for designing specific metalloproteinase inhibitors to intervene or eradicate breast tumor. Most importantly, the success of defining substrate specificity of MMP-2 and MMP-9 validates the feasibility of profiling protease activity via substrate phage display library technique.				
14. SUBJECT TERMS Breast Cancer				15. NUMBER OF PAGES 11
				16. PRICE CODE
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT Unlimited	

NSN 7540-01-280-5500

Standard Form 298 (Rev. 2-89)
Prescribed by ANSI Std. Z39-18
298-102

FOREWORD

Opinions, interpretations, conclusions and recommendations are those of the author and are not necessarily endorsed by the U.S. Army.

☒ Where copyrighted material is quoted, permission has been obtained to use such material.

☒ Where material from documents designated for limited distribution is quoted, permission has been obtained to use the material.

☒ Citations of commercial organizations and trade names in this report do not constitute an official Department of Army endorsement or approval of the products or services of these organizations.

N/A In conducting research using animals, the investigator(s) adhered to the "Guide for the Care and Use of Laboratory Animals," prepared by the Committee on Care and use of Laboratory Animals of the Institute of Laboratory Resources, national Research Council (NIH Publication No. 86-23, Revised 1985).

N/A For the protection of human subjects, the investigator(s) adhered to policies of applicable Federal Law 45 CFR 46.

N/A In conducting research utilizing recombinant DNA technology, the investigator(s) adhered to current guidelines promulgated by the National Institutes of Health.

N/A In the conduct of research utilizing recombinant DNA, the investigator(s) adhered to the NIH Guidelines for Research Involving Recombinant DNA Molecules.

N/A In the conduct of research involving hazardous organisms, the investigator(s) adhered to the CDC-NIH Guide for Biosafety in Microbiological and Biomedical Laboratories.


 6-23-00
PI - Signature Date

Table of Contents

Cover.....	1
SF 298.....	2
Foreword.....	3
Table of Contents.....	4
Introduction.....	5
Body.....	5-8
Key Research Accomplishments.....	9
Reportable Outcomes.....	9
Conclusions.....	9
References.....	10-11
Appendices.....	N/A

Introduction:

The purpose of my study is to obtain a collective knowledge of proteases which activity are differentially overexpressed during breast cancer progression using phage display technology. The profile of proteases or "protease fingerprint" will then be validated as a diagnostic tool for predicting breast cancer progression.

Body:

It is well-documented fact that early detection of breast tumors in patients can have significant impact on the survival rate of patients. Therefore, researchers have been actively searching for biomarkers which can be used to predict breast cancer progression. A good biomarker for breast cancer should have protein expression pattern correlates well with either the onset of malignant tumor or the tumor progression. In this fellowship proposal, I suggest that a profile of protease activity can be used as a diagnostic tool to predict breast cancer progression. Proteases capable of modifying extracellular matrix such as matrix metalloproteinase 2, 7 and 9 have been shown to involve in breast tumor metastasis, invasion, and angiogenesis (Ding, Coombs et al. 1995; Smith, Shi et al. 1995; Ke, Coombs et al. 1997; Coombs, Bergstrom et al. 1998). They are differentially overexpressed either in the stromal cells surrounding the tumor cells or in the tumor cells (Liotta 1992; Davies, Brown et al. 1993; Stetler-Stevenson, Liotta et al. 1993; Wang, Fu et al. 1994; DeClerck, Imren et al. 1997; Noel, Gilles et al. 1997). However, since these proteases can exist as three forms: inactive form with the attachment of the pro-domain, active form, inactive form complexed with endogenous inhibitors. Therefore, by monitoring the actual protease activity in tumor progression, we can grasp the alteration of tumor development and compile the information obtained into a predictable pattern for tumor progression. To approach this goal, I offered three aims in this proposal.

Aim#1: Construct a substrate phage display library to use as a tool for studying protease fingerprint.

It has been well established that the true assessment of the activity a protease is by measuring the hydrolysis of a peptide or protein substrate by the protease. Therefore, a substrate-based hydrolysis assay will be ideal for monitoring protease activity present during breast tumor development. Furthermore, several different proteases have been reported to involve in breast tumor progression and more proteases might yet to be discovered to involve in different processes of tumor progression, large amount of substrate sequences must be present in the assay to screen for protease activity in vivo to ensure a fair and unbiased assessment for proteases with different substrate preferences and perhaps novel proteases. In aim 1, I hypothesized that the substrate phage display library technique is an ideal assay that can be used to assess the activity of multiple proteases in vivo. Several substrate phage display libraries has been constructed by other researchers and used to extract information of preferred substrate sequences of disease related proteases such as plasminogen activators, thrombin, and MMP-7 (Ding, Coombs et al. 1995; Smith, Shi et al. 1995; Ke, Coombs et al. 1997; Coombs, Bergstrom et al. 1998). Moreover, large amount of randomized sequences can be obtained from a substrate display library ($> 1 \times 10^8$ sequences per library) to ensure a good representation of substrate sequences to detect the activity of multiple proteases.

To use as a tool of investigating protease activity, I construct a substrate phage display library which displays randomized hexameric inserts on the surface of phage. It is a modified version of polyvalent phage display system previously constructed by Scott and Smith (Smith 1985; Scott 1992; Smith 1993). The randomized hexameric sequences would mimic the peptide substrates and fit into the protease enzymatic pocket. In addition, I engineered a FLAG epitope tag at the N-terminal of the gIIIp-hexamer fusion protein to facilitate the separation of phages displayed preferred substrate sequences from other phages after each round of protease substrate selection. When a protease is exposed to the preferred substrate sequence displayed on the surface of phage, it will cleave the substrate sequence and liberate both the substrate and the FLAG epitope tag from the surface of phage. Subsequently, phages display protease preferred substrate sequences could be enriched by immuno-precipitation using a monoclonal anti-FLAG antibody and further amplified. A graphic illustration of this procedure can be found in figure 1 in the next page.




Figure 1. Scheme for selection of substrates by phage display technique.

Three substrate phage display libraries described above were constructed separately and represented by 1.8×10^8 , 2.4×10^8 , and 6.8×10^8 independent clones. I have confirmed the expression of FLAG epitope by Western Blot analysis, and a monoclonal antibody against the FLAG epitope can be used to immuno-precipitate phage. These libraries are at least as complex as any substrate phage libraries reported so far (Ding, Coombs et al. 1995; Smith, Shi et al. 1995; Ke, Coombs et al. 1997; Coombs, Bergstrom et al. 1998). This level of complexity is crucial to ensure 90% confidence level of including all possible combinations of randomized hexamer sequences. In which case, I am confident that these libraries can be used to explore substrates selectivity among different proteases.

Aim#2: Characterize profile of protease activity present in the blood of mice bearing non-metastatic breast cancer tumors and mice bearing metastatic breast cancer tumors.

Matrix metalloproteinases have been shown to play important roles in breast tumor invasion, metastasis, and angiogenesis (Liotta 1992; Stetler-Stevenson, Liotta et al. 1993; Noel, Gilles et al. 1997). Particularly, matrix metalloproteinase 2 and 9 have been studied extensively for their direct involvement in all of the processes required during tumor progression (Festuccia, Bologna et al. 1996; Sier, Kubben et al. 1996; Stearns and Stearns 1996; Stearns and Stearns 1996; Gohji, Fujimoto et al. 1998; Itoh, Ito et al. 1998; Kurizaki, Toi et al. 1998; Leppert, Ford et al. 1998; Moses, Wiederschain et al. 1998; Yu and Stamenkovic 2000). Like other matrix metalloproteinases, MMP-2 and MMP-9 are capable of cleaving extracellular matrix components such as collagen and laminin. Increased gene and protein expression of both proteases has been reported to correlate with tumor invasion and metastasis (Brooks, Stromblad et al. 1996; Festuccia, Bologna et al. 1996; Sier, Kubben et al. 1996; Stearns and Stearns 1996; Stearns and Stearns 1996; Gohji, Fujimoto et al. 1998; Kurizaki, Toi et al. 1998; Moses, Wiederschain et al. 1998). Also, from the immunohistological studies, MMP-2 and MMP-9 were found to localize in the tumor tissue (Brooks, Stromblad et al. 1996; Stearns and Stearns 1996; Stearns and Stearns 1996). To illustrate how these two proteases can be utilized by tumor cells, a mechanism of cell-surface association and activation for MMP-2 has been proposed to be used by tumor cells for invasion and metastasis (Strongin, Marmer et al. 1993; Strongin, Collier et al. 1995). As a validation of the substrate phage display library I constructed, I chose to define substrate specificity of MMP-2 and MMP-9 using purified proteins. Several advances could come from the study of substrate specificity. First, specific substrate sequences preferred by one protease over the other will help us distinguish functions of these two proteases. Unknown physiological substrates can potentially be revealed by defining the substrate preferences of each protease. Furthermore, the selected specific substrate sequences can serve as lead compounds for therapeutic and diagnostic development.

Principal Investigator: Chen, Emily

Since the hemopexin domain of both proteases have been shown to influence only the cell-surface association but not catalytic function of the protease (Murphy, Willenbrock et al. 1992), I chose to conduct the substrate specificity comparison with only the purified MMP-2 and MMP-9 catalytic domain. Embryonic kidney 293 cells were transfected separately with cDNA of either truncated MMP-2 or MMP-9 which includes only the pro-domain and truncated catalytic domain (no fibronectin-like domain) as described previously (Murphy, Willenbrock et al. 1992). Then, MMP-2 and MMP-9 protein were purified from conditioned media using gelatin affinity column as reported previously (Hipps, Hembry et al. 1991). The activity and purity of the purified MMP-2 and MMP-9 were verified by gelatin zymography and active site titration using a synthetic metalloproteinase inhibitor, ilomastat (Galardy, Cassabonne et al. 1994; Galardy, Grobelny et al. 1994). To select for MMP-2 substrates, 2×10^{10} phage from the initial library consisting of 2.4×10^8 independent clones were incubated with 10ug/ml of purified MMP-2 at 37°C.

Concurrently, a control reaction of 2×10^{10} phage with no protease was set up and incubated at the same condition as the selection reaction. Following the protease treatment, 100ug of monoclonal anti-FLAG antibody were added to both reactions, and the reactions were incubated at 4°C over night with rocking. Following the overnight incubation, pansorbin, which is inactivated bacterial cells expressing high concentration of protein A on the surface, was added to both reactions to precipitate the FLAG:IgG complexes, leaving the phage with cleaved substrate sequences in solution. The remaining phage, which carries MMP-2 preferred substrate sequences, was then amplified by infecting K91 bacteria and used for the subsequent round of selection.

For facilitate the screening of phage clones that carry protease preferred substrate sequences, a substrate phage ELISA was developed to rapidly identify desired phage clones. This assay has two major advantages. First, it eliminates the time consuming and tedious phage purification procedures by capturing the phage from the over night phage culture. Second, the readout of this assay yield a semi-quantitative indication of the rate of hydrolysis by the protease for individual sequences. Following two rounds of selection with purified MMP-2, I performed substrate phage ELISA to identify and characterize the ability of individual phage clones to be cleaved by MMP-2 as described in figure 2 below.

The substrate phage ELISA is carried out in 96-well microtiter plate format. A 96 well plate is coated with an anti-phage (gVIIIp) antibody. The wells are then incubated with 150ul supernatant from the over night culture of each tested phage clone. Four wells are coated with the same test phage clone: two wells as control with no protease, and two wells are for protease treatment. After two hours of phage binding at 4°C, 5ug/ml of MMP-2 were added to the designated wells and the plate is incubated at 37°C. In protease treatment reaction, MMP-2 will cleave the substrate and liberate the FLAG epitope from the phage. After protease incubation, the loss of FLAG epitope is detected by an anti-FLAG polyclonal antibody, followed by HRP-conjugated secondary antibody. Protease cleavage is indicated by the loss of FLAG epitope. This assay not only confirms that a peptide is a substrate, it also establishes a relative rate of hydrolysis between the substrates, that is the rate at which each substrate is hydrolyzed. In addition, it also allows me to test many potential substrates in one assay. Up to 100 substrates can easily be tested in one day.




Figure 2. Scheme for substrate phage ELISA.

Principal Investigator: Chen, Emily

A number of phage clones were selected by substrate phage ELISA with the loss of greater than 10% of the FLAG epitope due to the cleavage of substrate by MMP-2. DNA sequence analysis was performed for the selected phage clones to obtain the substrate sequences. Subsequent substrate sequence analysis revealed that three major classes of peptide substrates for MMP-2 were observed and summarized in table 1 in the next page. First major class represents a previously defined substrate preference for MMP-2 and MMP-9. Second and third classes represent two families of previously undefined substrates that could potentially distinguish MMP-2 from MMP-9 based on the data of substrate ELISA.

MMP-2 substrate classes	
Class 1: P-X-X-Hydrophobic amino acid	Previously defined substrates
Class 2: L-X-X-Hydrophobic amino acid	Novel substrates
Class 3: S-X-L	Novel undefined substrates

Table 1. Summary of three classes of MMP-2 peptide substrates.

One peptide from the PXX(hydrophobic amino acids) family, A-3, and one peptide from the LXX(hydrophobic amino acids), A49, were synthesized and their Kcat/Km values were determined for both MMP-2 and MMP-9. To determine the Kcat and Km value, I used a fluorescamine based assay(Fields, Van Wart et al. 1987). Fluorescamine is intrinsically nonfluorescent but reacts in milliseconds with primary aliphatic amines, including peptides and proteins, to yield a fluorescent derivative. After incubation of protease and peptide, the peptide is cleaved and the cleavage can be detected by addition of fluorescamine and read by a fluorescent microplate reader. Four different concentrations of each peptide were used to obtain the double reciprocal plot of initial velocity and substrate concentration. Km and Kcat values were then determined from the graph. The Kcat/Km values are listed in table 2 below.

Peptide	Sequence	MMP-2			MMP-9		
		Kcat/Km (M ⁻¹ S ⁻¹)	Kcat (S ⁻¹)	Km (mM)	Kcat/Km (M ⁻¹ S ⁻¹)	Kcat (S ⁻¹)	Km (mM)
A-3	S G A K P R A ↓ L	18414.6	3.02	0.164	39591.8	2.91	0.074
A49	S G L R L A A ↓ I	4208.3	1.01	0.24	270	*	*

Interestingly, the kinetic data of these two peptides confirmed the substrate ELISA result. This is first time that a MMP-2 specific peptide substrate sequence is identified. These MMP-2 specific substrate sequences could potentially be used as lead compounds for designing specific metalloproteinase inhibitors. Also, these unique sequences can be modified and then use as a tool to explore the physiological roles of MMP-2 and MMP-9 by discovering new physiological substrates in vivo. Furthermore, being able to define the substrate specificity of these two structurally and functionally related proteases would validate my hypothesis that the substrate phage display library can be used for profiling of different protease activity.

Future plans:

Synthesis of more MMP-2 specific peptides from both class 2 and class 3 MMP-2 substrate sequence is in progress. Their Kcat/Km values will be determined for both MMP-2 and MMP-9 to provide further confirmation of substrate specificity. Afterwards, I will launch to develop the method of protease profiling in vivo as described in aim#2. The mouse breast cancer model will be developed to obtain blood samples. Subsequently, I will select protease substrates using blood samples from normal mice, mice bearing non-metastatic tumor, and mice bearing metastatic tumor.

Principal Investigator: Chen, Emily

Key research accomplishments:

- I constructed three substrate phage display libraries with the highest complexity ever reported in the literature.
- I found new substrate sequences that are preferred by MMP-2 over MMP-9. These substrate sequences have therapeutic values because they can potentially be used as lead compounds for designing specific inhibitors.

Reportable outcomes:

These data were presented in the Annual Symposium of the Molecular Pathology Graduate Program of UCSD School of Medicine in June 22, 2000.

Conclusions/Implications:

- New families of MMP-2 substrates have been discovered. These substrates have selectivity for MMP-2 over MMP-9
- Substrate specificity appears to be governed by the S3 constituent of the substrate
- Our findings suggest many important substrates for MMP-2 may have been overlooked. These could clarify the biological role of MMP-2 (and MMP-9).

Reference:

- Brooks, P. C., S. Stromblad, et al. (1996). "Localization of matrix metalloproteinase MMP-2 to the surface of invasive cells by interaction with integrin alpha v beta 3." Cell **85**(5): 683-93.
- Coombs, G. S., R. C. Bergstrom, et al. (1998). "Substrate Specificity of Prostate-Specific Antigen (PSA)." Chemistry and Biology **5**(9): 475-488.
- Davies, B., P. D. Brown, et al. (1993). "A synthetic matrix metalloproteinase inhibitor decreases tumor burden and prolongs survival of mice bearing human ovarian carcinoma xenografts [published erratum appears in Cancer Res 1993 Aug 1;53(15):3652]." Cancer Research **53**: 2087-2091.
- DeClerck, Y. A., S. Imren, et al. (1997). "Proteases and protease inhibitors in tumor progression." Advances in Experimental Medicine & Biology **425**: 89-97.
- Ding, L., G. S. Coombs, et al. (1995). "Origins of the specificity of tissue-type plasminogen activator." Proc.Natl.Acad.Sci.U.S.A. **92**: 7627-7631.
- Festuccia, C., M. Bologna, et al. (1996). "Increased matrix metalloproteinase-9 secretion in short-term tissue cultures of prostatic tumor cells." International Journal of Cancer **69**(5): 386-93.
- Fields, G. B., H. E. Van Wart, et al. (1987). "Sequence specificity of human skin fibroblast collagenase. Evidence for the role of collagen structure in determining the collagenase cleavage site." Journal of Biological Chemistry **262**(13): 6221-6.
- Galardy, R. E., M. E. Cassabonne, et al. (1994). "Low molecular weight inhibitors in corneal ulceration." Annals of the New York Academy of Sciences **732**: 315-23.
- Galardy, R. E., D. Grobelny, et al. (1994). "Inhibition of angiogenesis by the matrix metalloprotease inhibitor N-[2R-2-(hydroxamidocarbonylmethyl)-4-methylpentanoyl]-L-tryptophan methylamide." Cancer Research **54**(17): 4715-8.
- Gohji, K., N. Fujimoto, et al. (1998). "Serum matrix metalloproteinase-2 and its density in men with prostate cancer as a new predictor of disease extension." International Journal of Cancer **79**(1): 96-101.
- Hipps, D. S., R. M. Hembry, et al. (1991). "Purification and characterization of human 72-kDa gelatinase (type IV collagenase). Use of immunolocalisation to demonstrate the non-coordinate regulation of the 72-kDa and 95-kDa gelatinases by human fibroblasts." Biological Chemistry Hoppe-Seyler **372**(4): 287-96.
- Itoh, Y., A. Ito, et al. (1998). "Plasma membrane-bound tissue inhibitor of metalloproteinases (TIMP)-2 specifically inhibits matrix metalloproteinase 2 (gelatinase A) activated on the cell surface." Journal of Biological Chemistry **273**(38): 24360-7.
- Ke, S. H., G. S. Coombs, et al. (1997). "Distinguishing the Specificities of Closely Related Proteases." The Journal of Biological Chemistry **272**(26): 16603-16609.
- Kurizaki, T., M. Toi, et al. (1998). "Relationship between matrix metalloproteinase expression and tumor angiogenesis in human breast carcinoma." Oncology Reports **5**(3): 673-7.
- Leppert, D., J. Ford, et al. (1998). "Matrix metalloproteinase-9 (gelatinase B) is selectively elevated in CSF during relapses and stable phases of multiple sclerosis." Brain **121**(Pt 12): 2327-34.
- Liotta, L. A. (1992). "Cancer cell invasion and metastasis." Sci.Am. **266**: 54-9, 62-3.
- Moses, M. A., D. Wiederschain, et al. (1998). "Increased incidence of matrix metalloproteinases in urine of cancer patients." Cancer Research **58**(7): 1395-9.
- Murphy, G., F. Willenbrock, et al. (1992). "The C-terminal domain of 72 kDa gelatinase A is not required for catalysis, but is essential for membrane activation and modulates interactions with tissue inhibitors of metalloproteinases [published erratum appears in Biochem J 1992 Jun 15;284(Pt 3):935]." Biochem J **283**(Pt 3): 637-41.
- Noel, A., C. Gilles, et al. (1997). "Emerging roles for proteinases in cancer." Invasion & Metastasis **17**(5): 221-39.
- Scott, J. K. (1992). "Discovering peptide ligands using epitope libraries." Trends in Biochemical Sciences **17**: 241-245.

Principal Investigator: Chen, Emily

- Sier, C. F., F. J. Kubben, et al. (1996). "Tissue levels of matrix metalloproteinases MMP-2 and MMP-9 are related to the overall survival of patients with gastric carcinoma." British Journal of Cancer **74**(3): 413-7.
- Smith, G. P. (1985). "Filamentous fusion phage: novel expression vectors that display cloned antigens on the virion surface." Science **228**: 1315-1317.
- Smith, G. P. (1993). "Surface display and peptide libraries." Gene **128**: 1-2.
- Smith, M. M., L. Shi, et al. (1995). "Rapid identification of highly active and selective substrates for stromelysin and matrilysin using bacteriophage peptide display libraries." J Biol Chem **270**(12): 6440-9.
- Stearns, M. and M. E. Stearns (1996). "Evidence for increased activated metalloproteinase 2 (MMP-2a) expression associated with human prostate cancer progression." Oncology Research **8**(2): 69-75.
- Stearns, M. E. and M. Stearns (1996). "Immunohistochemical studies of activated matrix metalloproteinase-2 (MMP-2a) expression in human prostate cancer." Oncology Research **8**(2): 63-7.
- Stetler-Stevenson, W. G., L. A. Liotta, et al. (1993). "Extracellular matrix 6: role of matrix metalloproteinases in tumor invasion and metastasis." FASEB.J. **7**: 1434-1441.
- Strongin, A. Y., I. Collier, et al. (1995). "Mechanism of cell surface activation of 72-kDa type IV collagenase. Isolation of the activated form of the membrane metalloprotease." J Biol Chem **270**(10): 5331-8.
- Strongin, A. Y., B. L. Marmer, et al. (1993). "Plasma membrane-dependent activation of the 72-kDa type IV collagenase is prevented by complex formation with TIMP-2." J. Biol. Chem. **268**(19): 14033-9.
- Wang, X., X. Fu, et al. (1994). "Matrix metalloproteinase inhibitor BB-94 (batimastat) inhibits human colon tumor growth and spread in a patient-like orthotopic model in nude mice." Cancer Research **54**: 4726-4728.
- Yu, Q. and I. Stamenkovic (2000). "Cell surface-localized matrix metalloproteinase-9 proteolytically activates TGF-beta and promotes tumor invasion and angiogenesis." Genes & Development **14**(2): 163-76.